

Regularized Kerr-Newman Solution as a Gravitating Soliton

Alexander Burinskii

*Theoretical Physics Laboratory, NSI, Russian Academy of Sciences, B. Tulkaya 52 Moscow 115191 Russia**

The charged, spinning and gravitating soliton is realized as a regular solution of the Kerr-Newman field coupled with a chiral Higgs model. A regular core of the solution is formed by a domain wall bubble interpolating between the external Kerr-Newman solution and a flat superconducting interior. An internal electromagnetic (em) field is expelled to the boundary of the bubble by the Higgs field. The solution reveals two new peculiarities: (i) the Higgs field is oscillating, similar to the known oscillon models, (ii) the em field forms on the edge of the bubble a Wilson loop, resulting in quantization of the total angular momentum.

PACS numbers: 11.27.+d, 04.20.Jb, 04.70.Bw

1. Introduction.

There are many pieces of evidence that black holes (BHs) may be related with elementary particles [1]. Being regularized, BHs turn out to be similar to gravitating solitons [2, 3] which are considered as regular field models of the extended particles in the electroweak sector of the standard model [2, 4–6]. The Kerr-Newman (KN) solution plays a fundamental role as a model of a rotating BH and as a convenient laboratory for the study of consistency between quantum theory and gravity [7–9]. The aim of this paper is to obtain a regular version of the KN solution which may also be considered as a field model of spinning particle compatible with general relativity.

The attempts to incorporate spin and gravity in the soliton models encounter the problems related with choosing appropriate field models. As a result, the models turn out to be difficult for analysis and admit only numerical solutions [3, 10, 11]. In this paper we use an alternative approach starting from a gravitational side – specific structure of the KN solution. Our previous attempts to obtain its regular version [12, 13] were based on the bag-like Higgs model [12] and the Gürses and Gürsey (GG) ansatz for the regular Kerr-Schild (KS) metrics [13, 14]. However, there were problems of consistency in the electromagnetic (em) sector, and subsequent attention was drawn to the models based on non-linear electrodynamics (NED), where the existence of the regular charged non-rotating BH solutions was explicitly demonstrated (in particular in [15, 16]). Nevertheless, the investigation of the corresponding *rotating* solutions led to contradicting results.[44] In the present treatment, we return to the bag-like model considered in [12] based on the Higgs mechanism of broken symmetry and show that the problems in the em sector of the bag-like Higgs model may be successfully resolved, leading to new important physical results.

It was obtained by Carter in 1968 that the KN solution has $g = 2$ as that of the Dirac electron [19], and there followed a series of the works on the KN solution as a model of a spinning particle, in particular [12, 18–27]. Since the spin of an electron is very high, the BH horizons of the KN solution disappear, opening a naked singular ring which should be replaced by a regular source. We consider here a regular field model of the KN source, in which the KN solution is modified near singular ring and replaced by a source forming the bubble which has a superconducting internal vacuum state formed by the Higgs mechanism of the broken symmetry. The Higgs sector of our field model is similar to that of the well-known Nielsen-Olesen (NO) model for a regular vortex in superconducting media [28]. However, to describe a domain wall interpolating between the exact external KN solution and the flat superconducting interior, we need the Higgs model containing a few chiral fields. We found that the regular internal solution is unambiguously determined by the external KN field, and the regularized KN solution acquires two important peculiarities:

- (i) the internal Higgs field turns out to be spinning and oscillating with frequency $2m$, similar to the models of oscillating Q-balls [29, 30] and bosonic stars [31].
- (ii) the external em field acquires on the edge of the bubble a closed Wilson loop, which entails quantization of angular momentum.

2. The Kerr singular ring, twosheetedness and the bubble source.

The problem of regular source of the KN solution becomes complicated by twosheetedness of the Kerr geometry. The KS form of metric $g_{\mu\nu} = \eta_{\mu\nu} + 2Hk_\mu k_\nu$ is coupled to the auxiliary Minkowski metric $\eta^{\mu\nu}$ with the Cartesian

* E-mail address: bur@ibrae.ac.ru

coordinates $x^\mu = (t, x, y, z)$. For the KN solution function H has the form

$$H = \frac{mr - e^2/2}{r^2 + a^2 \cos^2 \theta}, \quad (1)$$

where Kerr's oblate spheroidal coordinates r, θ and ϕ_K are related with the Cartesian coordinates as follows

$$x + iy = (r + ia) \exp\{i\phi_K\} \sin \theta, \quad z = r \cos \theta. \quad (2)$$

The coordinate r covers the Kerr space-time twice, for $r > 0$ and for $r < 0$, forming the 'positive' and 'negative' sheets which are analytically connected via disk $r = 0$. On the boundary of the disk, $r = \theta = 0$, function H is singular, which corresponds to the Kerr singular ring.[45] To remove the Kerr singularity and twosheetedness, they have to be covered by a material source. Israel suggested to truncate the negative KN sheet along the disk spanned by the Kerr ring and considered this disk, coupled with the Kerr ring, as a model of the KN source [20]. This source was consistently matched with KN exterior but resulted in the very exotic superluminal matter of the disk. Then Hamity showed in [32] that the disk has to be relativistically rotating and consisting of the matter with zero energy density and negative pressure which diverge near the Kerr singular ring. Finally, López [22] determined the form of the source, matching continuously the KN exterior with the bubble determined by the oblate coordinate $r = r_0 = e^2/2m$. The function H vanishes at this r_0 which allowed him to match exterior continuously with the flat interior. [46] This model was consistent, however, like other shell-like models, it needed the Poincaré stress compensating the repulsion of the charges distributed over the shell. Meanwhile, the necessary tangential stress appears naturally in the domain wall field models, and we exploit it in the presented model of the KN source.

3. Field model.

We consider for the matter fields the Abelian Higgs model with the Lagrangian [28]

$$\mathcal{L}_{NO} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}(\mathcal{D}_\mu\Phi)(\mathcal{D}^\mu\Phi)^* + V(r), \quad (3)$$

coupled with gravity, so $\mathcal{D}_\mu = \nabla_\mu + ieA_\mu$ are to be covariant derivatives. To form a bubble source we need a domain wall potential $V(r)$ interpolating smoothly between the external vacuum state, $V^{ext} = 0$, and the internal 'false' vacuum with $V^{int} = 0$. The external vacuum should yield the longrange external em field of the KN solution, while the interior of the bubble should be superconducting, leading to a shorrange em field to provide regularization. It is opposite to the structure of the NO model, but corresponds to the structure of Witten's field model for superconducting cosmic strings [34], and we use the supersymmetric version of this model suggested by Morris [35] with three chiral fields $\Phi^{(i)} = \{\Phi, Z, \Sigma\}$, $i = 1, 2, 3$, controlled by the Lagrangian

$$\mathcal{L}_{matter} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}\sum_i(\mathcal{D}_\mu^{(i)}\Phi^{(i)})(\mathcal{D}^{(i)\mu}\Phi^{(i)})^* + V, \quad (4)$$

where the auxiliary field Z is real and uncharged. We need only one gauge field A_μ associated with the chiral field $\Phi^{(1)} \equiv \Phi$; therefore, $F_{\mu\nu} = A_{\mu,\nu} - A_{\nu,\mu}$ and the corresponding covariant derivations will be $\mathcal{D}_\mu^{(1)} = \nabla_\mu + ieA_\mu$, $\mathcal{D}_\mu^{(2)} = \mathcal{D}_\mu^{(3)} = \nabla_\mu$.

The potential V is determined via the superpotential W by the relation $V(r) = \sum_i |\partial_i W|^2$, where $\partial_1 = \partial_\Phi$, $\partial_2 = \partial_Z$, $\partial_3 = \partial_\Sigma$. The given by Morris superpotential is

$$W = \lambda Z(\Sigma\bar{\Sigma} - \eta^2) + (cZ + \mu)\Phi\bar{\Phi}, \quad (5)$$

where c, μ, η, λ are real constants. This potential produces a domain wall interpolating between the internal $V^{(int)} = 0$ and external vacuum states $V^{(ext)} = 0$, determined by the condition $\partial_i W = 0$. We use the thin wall approximation, assuming that the depth of the wall ξ , is much smaller than its position r_0 , which yields

- (in) for $r < r_0$: $Z = -\mu/c$; $\Sigma = 0$; $|\Phi| = \eta\sqrt{\lambda}/c$,
- (out) for $r > r_0 + \xi$: $Z = 0$; $\Phi = 0$; $\Sigma = \eta$.

In the external region we have $V^{(ext)} = 0$, and the unique nonzero chiral field Σ is constant. There survives only electromagnetic matter, which results in the Einstein-Maxwell equations leading to the usual KN solution.

For the interior, $r < r_0$, we have $V^{(int)} = 0$, and the unique nonzero chiral field is Φ . As a result, the Lagrangian (4) is reduced to (3), leading to the usual equations for the coupled with gravity Maxwell-Higgs system

$$\mathcal{D}_\nu^{(1)}\mathcal{D}^{(1)\nu}\Phi = \partial_\Phi V = 2\Phi[c^2|\Phi|^2 - \lambda c\eta^2] = 0, \quad (6)$$

$$\nabla_\nu\nabla^\nu A_\mu = I_\mu = \frac{1}{2}e|\Phi|^2(\chi_{,\mu} + eA_\mu). \quad (7)$$

However, as we see in the next section, there is a strong reason to choose parameters of the model which lead to the flat space inside the bubble. As a result, these equations are simplified, allowing us to obtain exact solutions.

4. Regularization of the KN metric.

The KS metrics which describe a smooth interpolation from the external metric of the KN solution to a regular internal space-time where considered in [12, 13] on the base of the GG ansatz [14]. For the external region one uses the exact KN solution in the KS form

$$g_{\mu\nu} = \eta_{\mu\nu} + 2Hk_\mu k_\nu, \quad (8)$$

with $H = \frac{mr - e^2/2}{r^2 + a^2 \cos^2 \theta}$. The KN electromagnetic (em) field is given by the vector potential

$$A_{KN}^\mu = Re \frac{e}{r + ia \cos \theta} k^\mu, \quad (9)$$

where $k^\mu(x^\mu)$ is the null vector field tangent to a vortex of the null geodesic lines, the Kerr principal null congruence (PNC). The field $k^\mu(x)$ has the form [19]

$$k_\mu dx^\mu = dr - dt - a \sin^2 \theta d\phi_K. \quad (10)$$

For metric inside the bubble, $r < r_0$, one uses the KS form (8) with the function

$$H = f(r)/\Sigma, \quad \Sigma = (r^2 + a^2 \cos^2 \theta). \quad (11)$$

It describes the external KN metric for $f(r) = f_{KN} = mr - e^2/2$, and any smooth function $f(r)$ interpolating between $f_{KN}(r)$ and $f_{int}(r)$ provides a smooth transfer to a metric inside the bubble. If we set for the interior $f(r) = f_{int} = \alpha r^4$, the Kerr singularity will be suppressed. The GG form of function H (11) allows one to describe in a universal manner the de Sitter and Anti-de Sitter solutions, and match smoothly the external and internal metrics of different types. Besides, it demonstrates a correspondence between rotating and non-rotating Kerr-Schild solutions which simplifies analysis of the rotating solutions. The Kerr space-time is foliated into ellipsoidal layers $r = \text{const.}$ which are rigidly rotating with angular velocities depending on the Kerr oblate coordinate r , $\Omega(r) = \frac{a}{a^2 + r^2}$. All the fields on the layers may be transformed to the co-rotating coordinate systems by the corresponding local Lorentz transformation, where they take a simple diagonal form [12–14, 36])

$$T_{\mu\nu} = \text{diag}(\rho, p_{tan}, p_{tan}, p_{rad}), \quad (12)$$

where

$$\rho = -p_{rad} = \frac{1}{4\pi}(f'r - f)/\Sigma^2, \quad p_{tan} = \rho - \frac{1}{8\pi}f''/\Sigma, \quad (13)$$

corresponding to the non-rotating solutions. Because of that many results obtained for the non-rotating case ($a = 0$ and $\Sigma = r^2$ in (11)), may easily be translated back to the rotating case by the replacement $\Sigma \rightarrow r^2 + a^2 \cos^2 \theta$. In particular, it allows one to match smoothly the external and internal metrics with rotation.

Setting $f_{int} = \alpha r^4$, one obtains from (12) and (13) that it corresponds a regular rotating internal space-time of the constant curvature $R = -24\alpha$, [12, 13]. The energy density inside the bubble will be $\rho = \frac{3\alpha}{4\pi}$, which generates de Sitter interior for $\alpha > 0$, anti de Sitter interior for $\alpha < 0$ and the flat interior corresponding to the López model for $\alpha = 0$. The three corresponding branches for the interior are shown in fig.1 (taken from [36]). They intersect the plot $f_{KN}(r)$ at the points $r_0 = r_{AdS}$, $r_0 = r_{dS}$ and $r_0 = r_{flat}$ correspondingly, showing the point where the external KN metric matches continuously with the interior. Selecting a thin layer of phase transition, $r_0 < r < r_0 + \xi$, one can take a smooth interpolating function $f(r)$ and provide smoothness of the general metric. In the thin domain wall limit $\xi \rightarrow 0$, the point of intersection $f_{int}(r) = f_{KN}(r)$ determines r_0 graphically. One sees that r_0 is solution of the equation $\alpha r_0^4 = mr_0 - e^2/2$, and since $\rho = \frac{3\alpha}{4\pi}$, one obtains for ADM mass the relation

$$m = e^2/2r_0 + \rho \frac{4}{3}\pi r_0^3, \quad (14)$$

where the first term on r.h.s. is the mass of the external em field for the charged sphere, and the second term is the mass inside this sphere with the matter density ρ . For small ξ the contribution of the ξ -shell to the mass m is negligible, and we obtain an important result that *the point of the phase transition r_0 is determined by the 'balance*




FIG. 1: Matching of the metric for regular bubble interior with metric of external KN field.

matter equation' (14). This analysis is extended to the rotating case, in which r has to be considered as the Kerr oblate ellipsoidal coordinate, and the boundary of bubble takes the form of an oblate rotating disk, [12, 13, 36].

We restrict further the treatment by the case of a flat interior corresponding to $\alpha = \rho = f(r) = 0$, $r < r_0$.

Stringy effect. In the rotating case the Arnowitt-Deser-Misner (ADM) mass of the KN source is determined by T_0^0 component of the stress-energy tensor in the asymptotically flat coordinate system. Contributions to the ADM mass coming from the interior of the bubble, the intermediate region, and the external electromagnetic field were calculated in [13]. For the flat interior, $\alpha = 0$, and an infinitely thin transitional shell we have: $m_{ADM}^{(int)} = 0$; and

$$m_{ADM}^{(shell)} = \frac{m}{2} \left[1 - \left(\frac{r_0}{a} + \frac{a}{r_0} \right) \arctan\left(\frac{a}{r_0}\right) \right]; \quad (15)$$

$$m_{ADM}^{(ext)} = \frac{m}{2} \left[1 + \left(\frac{r_0}{a} + \frac{a}{r_0} \right) \arctan\left(\frac{a}{r_0}\right) \right]. \quad (16)$$

For $a \gg r_0$, the second term in the square brackets of (16) tends to the expression $\delta^{string} \approx \frac{a\pi}{2r_0}$ which may be interpreted as a stringy contribution to the mass-energy caused by concentration of the electromagnetic field on the edge of bubble. [47] Stringy structure of the region near the Kerr ring was discussed many times, [23, 24, 37, 38]. Assuming that the closed Kerr string has a tension T with the rising potential $E \equiv m = Ta$, one can combine it with the basic KN relation $J = ma$, and obtain the linear Regge trajectory $J = \frac{1}{T}m^2$ with the slope $1/T$. However, one sees from (15) and (16) that the 'stringy' contributions from the shell and external em field are mutually cancelled, and the total mass $m_{ADM}^{(total)} = m_{ADM}^{(int)} + m_{ADM}^{(shell)} + m_{ADM}^{(ext)}$ turns out to be equal to m . Indeed, this result could be predicted *a priori*, since the total ADM mass is determined *only* by the asymptotical gravitational field, i.e. by the parameter m in function H , (1). The treatment of the Tolman mass expression, or the transitional zone of a finite thickness does not change the result. The partial stringy contributions to the mass are very intense ($\delta m/m > 100$), but mutual cancelling of the negative and positive contributions prevents exhibition of the stringy effect in the considered isolated system. However, the balance of matter may apparently be destroyed by an external field, and a considerable effect may be expected in the bound systems or in consequence of the radiative corrections.

5. Regularization of the KN electromagnetic field.

We assume that for $r < r_0$ the phase transition is completed and inside the bubble the field $\Phi(x)$ has the form $\Phi(x) = |\Phi(x)|e^{i\chi(x)}$ with a nonzero vev, $|\Phi(x)|_{r < r_0} = \Phi_0$. We have to obtain a regular solution of equation (7) for $r < r_0$ in the presumption of the flat interior, $\alpha = 0$, which fixes the boundary of bubble at $r_0 = r_e = e^2/2m$. The flat interior allows us to use the flat d'Alembertian and $\mathcal{D}_\mu = \partial_\mu + ieA_\mu$ in (7) and yields

$$\square A_\mu = I_\mu = e|\Phi|^2(\chi_{,\mu} + eA_\mu). \quad (17)$$

The current has to be expelled from the bulk of the superconducting interior to the boundary of the bubble and we should set in the interior $I_\mu = 0$, which yields

$$\square A_\mu^{(in)} = 0 = e|\Phi|^2(\chi_{,\mu} + eA_\mu^{(in)}). \quad (18)$$

The external KN field A_μ is given by (9) and (10). Its matching with the interior turns out to be nontrivial, since the Kerr angular coordinate ϕ_K (2) is very specific, and inside the bubble it turns out to be inconsistent with the

simple angular coordinate of the Higgs field, $\phi = -i \ln[(x + iy)/\rho]$, where $\rho = (x^2 + y^2)^{1/2}$. Using the relation between the corresponding differentials

$$d\phi_K = \phi + \frac{adr}{r^2 + a^2}, \quad (19)$$

we transform the KN vector potential on the boundary and inside the bubble to the form

$$A_\mu dx^\mu = \frac{-er}{r^2 + a^2 \cos^2 \theta} [dt + a \sin^2 \theta d\phi] + \frac{2erdr}{(r^2 + a^2)} \quad (20)$$

which shows that the radial component A_r is a perfect differential. On the external side of the bubble, $r = r_e + 0 = e^2/2m$, the value of the potential is

$$A_\mu dx^\mu|_{r_e+0} = \frac{-er_e}{r_e^2 + a^2 \cos^2 \theta} [dt + a \sin^2 \theta d\phi] + \frac{2er_e dr}{(r_e^2 + a^2)}. \quad (21)$$

The lines of the KN vector-potential in the equatorial plane, $\cos \theta = 0$, are tangent to the Kerr singular ring, [23, 24], and approaching the string-like edge of the KN bubble, the potential takes the form

$$A_\mu^{(str)} dx^\mu = -\frac{2m}{e} [dt + ad\phi] + \frac{2er_e dr}{(r_e^2 + a^2)}. \quad (22)$$

The tangent component on the edge takes the value $A_\phi^{(str)} = -2ma/e$. In agreement with (17) and $I_\mu = 0$, the component $A_\phi^{(str)}$ has to be matched with angular periodicity of the internal Higgs field $\Phi = \Phi_0 \exp(i\chi)$, which determines its ϕ -dependence, $\Phi \sim \exp\{in\phi\}$, n -integer. Therefore, the condition $I_\mu = 0$ demands mutual compensation of the gauge and Higgs field inside the bubble and leads to the closed Wilson loop on its edge. The integral

$$S = \oint e A_\phi^{(str)} d\phi = -4\pi ma \quad (23)$$

has to be matched with incursion of the Higgs phase $2\pi n$, which, due to the KN relation $J = ma$, entails $J = n/2$, where n is integer. Therefore, the consistent matching selects quantum values of the KN angular momentum.

For the time-like component A_0 inside the bubble, the condition $I_\mu = 0$ yields $\chi_{,0} = -eA_0^{(in)}(r)$, which determines $\chi_{,0}$ as a constant corresponding to the frequency of oscillation of the Higgs field, $\chi_{,0} = \omega = -eA_0^{(str)} = 2m$. Therefore, the vector potential inside the bubble has the components

$$A_0^{(in)} = -\frac{2m}{e}; \quad A_\phi^{(in)} = -\frac{2ma}{e}; \quad A_r^{(in)} = \frac{2erdr}{(r^2 + a^2)}. \quad (24)$$

This potential satisfies the left-hand side of equation (18), so far as it is gradient of the scalar function $\chi(t, r, \theta)$. At the same time, it also satisfies the right-hand side of (18), since it is compensated by the phase of the Higgs field.

The internal Higgs field takes the form

$$\Phi(x) = \Phi_0 \exp\{i\omega t - i \ln(r^2 + a^2) + 2in\phi\}. \quad (25)$$

For exclusion of the region of the closed loop at the equator, the time and ϕ components of the obtained vector field have a chock, crossing the boundary of the bubble, which generates circular currents distributed over the surface layer.

6. Consistency of the chiral model with gravity.

The complete stress-energy tensor may be decomposed into pure em part and contributions of the chiral fields

$$T_{\mu\nu}^{(tot)} = T_{\mu\nu}^{(em)} + \delta_{i\bar{j}} (\mathcal{D}_\mu^{(i)} \Phi^i) (\overline{\mathcal{D}_\nu^{(j)} \Phi^j}) - \frac{1}{2} g_{\mu\nu} [\delta_{i\bar{j}} (\mathcal{D}_\lambda^{(i)} \Phi^i) (\overline{\mathcal{D}^{(j)\lambda} \Phi^j}) + V], \quad (26)$$

In the exterior we have $V^{(ext)} = 0$, and the unique nonzero chiral field Σ tends to constant. Therefore, all the derivatives $\mathcal{D}_\mu^{(i)} \Phi^{(i)}$ will vanish, and $T_{\mu\nu}^{(tot)} = T_{\mu\nu}^{(em)}$, which leads to consistency with the Einstein-Maxwell equations and to the external KN solution. For the interior, $r < r_0$, we have the flat space, $\alpha = 0$, and $V^{(int)} = 0$, which

yields the unique nonconstant Higgs field $\Phi(x) = |\Phi(x)|e^{i\chi(x)}$. The Lagrangian (4) reduces to (3), leading to the equations (7). For the case of flat interior these equations were solved explicitly in sec.5. As a result, the stress-energy tensor (26) turns out to be consistent with the flat interior, and with the external KN metric. At the same time, its consistency in the zone of the phase transition is not established at this stage, and we may consider the obtained solution only as a thin-wall approximation.

7. Beyond the thin wall approximation.

The structure of the zone of the phase transition is very nontrivial and deserves especial study. In this section, we give only a preliminary treatment of the encountered problems. A thin but finite zone of the phase transition may be modelled by a planar vacuum domain wall, the stress-energy tensor of which has the following typical structure [40, 41]

$$T_{\mu\nu}^{DW} = \rho \text{diag}(1, -1, -1, 0), \quad (27)$$

indicating that the wall tension is equal to energy density. The stress-energy tensor of the considered chiral field model also has this form in the case of the planar vacuum domain wall without the em field. However, subtracting from (12) and (13) the contributions from the em field, we obtain the form

$$T_{\mu\nu} = |p_{tan}| \text{diag}(0, -1, -1, 0), \quad (28)$$

which differs from the typical ‘vacuum’ domain wall, indicating the presence of an unaccounted structure. One of the neglected phenomena is related to the penetration of the em field in the superconducting region, the zone of ‘penetration length’ [28], where the gauge field acquires a mass via the Higgs effect. In our case this zone forms an extra layer $U_\delta = [r_0 - \delta < r \leq r_0]$ on the boundary of the disk-like bubble. The massive gauge field concentrates on the edge of bubble in the equatorial plane, and forms a circular stringy region positioned very close to the former Kerr singular ring (the Cartesian distance from the Kerr ring is about ae^4). On the external side of the bubble, $r = r_0 + 0$, this massive field transfers into the gauge field of the above Wilson loop, while on the internal side of the bubble it creates circular currents, which are determined by the deviation of the real em field from the background solutions (24), (25) obtained in section 5. Substituting $A_\mu^{(\delta)} = A_\mu - A_\mu^{(in)}$ into (17) one obtains the equation

$$\square A_\mu^{(\delta)} = I_\mu = e^2 |\Phi|^2 A_\mu^{(\delta)} \quad (29)$$

which describes this effect.

The circular stringy region rotates together with the bubble with angular velocity $\Omega = ca/(a^2 + r_0^2)$. So, its linear velocity on the edge of bubble is $v_0 = c/\sqrt{1 + (r_0/a)^2} \sim c$. In particular, for the parameters of an electron $r_0/a = 137^{-1}$, and the linear velocity of the edge $v_0 \approx 0.9999c$ is almost lightlike. Therefore, we again arrive at the stringy structure which appears in the transition region. This structure is very nontrivial and important, and we intend to consider it in a separate paper.

8. Conclusion.

In the thin-wall approximation, we obtained the regular solution of the KN geometry which may be considered as a charged, spinning and gravitating soliton. The Higgs field inside its regular core is coherently oscillating, which reproduces the typical property of the oscillon models: spinning Q-balls [29, 30] and bosonic star models [31].

One of the new important peculiarities of this solution is the appearance of the closed Wilson loop at the boundary of the bubble source which leads to quantization of the total angular momentum. The closed vortexes were also obtained in some other soliton models [3, 29], but in our case the Wilson loop has a specific origin, being determined by the vortex structure of the Kerr singular ring.

The considered model is Abelian and may be related to the light spinning particles. Although, it may apparently be generalized to the non-Abelian case too. For the parameters of an electron, the considered regular solution predicts a disk-like core formed by the Higgs field, which is bounded by a closed circular current of the Compton radius. In principle, it agrees with the size of a dressed electron predicted by QED, but suggests some extra features of the spinning solutions consistent with general relativity. First is that the core is coherently oscillating and therefore, it should be considered as an integral element of the spinning particle. Second is the disk-like form of the core, which should apparently have experimentally observable consequences.[48]

Finally, we should note that the mysterious problem of twosheetedness of the KN spacetime is turned into its advantage, since the KN source forms a holographic structure [9] which is necessary for quantum treatments [7, 42]. The inner flat vacuum state may be extended analytically to the negative KN sheet which takes the role of a flat

in-vacuum space, separated from the physical out-sheet by the holographically dual boundary of the bubble, forming the bulk/boundary correspondence.

Acknowledgements.

The author thanks T.M. Nieuwenhuizen for hospitality in ITP of Amsterdam University and for discussions which stimulated reconsideration of this problem, and also thanks D. Stevens for conversations and useful reference.

-
- [1] G. 't Hooft, Nucl. Phys. B **335**, 138 (1990);
C. F. E. Holzhey and F. Wilczek, Nucl. Phys. B **380**, 447 (1992) [arXiv:hep-th/9202014].
A. Sen, Mod. Phys. Lett. A **10**, 2081 (1995) [arXiv:hep-th/9504147].
H. Nishino, Phys. Lett. B **359**, 77 (1995) [arXiv:hep-th/9504142].
 - [2] E. Radu and M. S. Volkov, "Existence of stationary, non-radiating ring solitons in field theory: knots Phys. Rept. **468**, 101 (2008) [arXiv:0804.1357 [hep-th]].
 - [3] B. Kleihaus, J. Kunz and Y. Shnir, Phys. Rev. D **70**, 065010 (2004) [arXiv:hep-th/0405169].
J. Kunz, U. Neemann, Ya. Shnir, Phys. Rev. D **75**, 125008 (2007).
 - [4] A. Kusenko, Phys. Lett. B **404**, 285 (1997).
 - [5] N. S. Manton, Phys. Rev. D **28**, 2019 (1983).
 - [6] R. F. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. D **10**, 4138 (1974).
 - [7] C. R. Stephens, G. 't Hooft and B. F. Whiting, Class. Quant. Grav. **11**, 621 (1994) [arXiv:gr-qc/9310006].
 - [8] A. Burinskii, Gen. Rel. Grav. **41**, 2281 (2009) [arXiv:0903.3162 [gr-qc]].
Reprinted in A. Burinskii, Int. J. Mod. Phys. D **18** 2351 (2009) [arXiv: 0903.3162,]
A. Burinskii, J. Phys. Conf. Ser. **222**, 012044 (2010) [arXiv:1001.0332 [gr-qc]].
 - [9] A. Burinskii, Theor. Math. Phys. **163**(3) 782 (2010), [arXiv:0911.5499].
 - [10] Y. Brihaye and B. Hartmann, Phys. Rev. D **79**, 064013 (2009).
 - [11] E. Seidel and W.-M. Suen, Phys. Rev. Lett. **66**, 1659 (1991).
 - [12] A. Y. Burinskii, Grav. Cosmol. **8**, 261 (2002) [arXiv:hep-th/0008129].
A. Burinskii, *Supersymmetric Bag Model as a Development of the Witten Superconducting String*, [arXiv:hep-th/0110011].
 - [13] A. Burinskii, E. Elizalde, S. R. Hildebrandt and G. Magli, Phys. Rev. D **65**, 064039 (2002) [arXiv:gr-qc/0109085].
 - [14] M. Gürses and F. Gürsey, J. Math. Phys. **16**, 2385 (1975).
 - [15] A. Burinskii and S. R. Hildebrandt, Phys. Rev. D **65**, 104017 (2002).
 - [16] I. Dymnikova, Class. Quant. Grav. **19**, 725 (2002).
 - [17] A. Burinskii and S. R. Hildebrandt, Grav. Cosmol. **9** 20 (2003).
 - [18] I. Dymnikova, Phys. Lett. B **639**, 368 (2006) [arXiv:hep-th/0607174].
 - [19] G. C. Debney, R. P. Kerr and A. Schild, J. Math. Phys. **10**, 1842 (1969).
 - [20] W. Israel, Phys. Rev. D **2**, 641 (1970).
 - [21] A.Ya. Burinskii, "Microgeons with spin," Sov. Phys. JETP **39**, 193 (1974), (Translated from Zh. Eksp. Teor. Fiz. , v. 66, no. 2, pp. 406-411, 1974.)
A.Ya. Burinskii, "Microgeons with Kerr metric," Russ. Phys. J. **17**, 1068 (1974), (doi: 10.1007/BF00901591),
[http://www.springerlink.com/content/n406x02205417878/]
 - [22] C. A. Lopez, Phys. Rev. D **30**, 313 (1984).
 - [23] A. Burinskii, Phys. Rev. D **52**, 5826 (1995) [arXiv:hep-th/9504139].
 - [24] A. Burinskii, Phys. Rev. D **70**, 086006 (2004) [arXiv:hep-th/0406063].
 - [25] H. I. Arcos and J. G. Pereira, Gen. Rel. Grav. **36**, 2441 (2004) [arXiv:hep-th/0210103].
 - [26] Th.M. Nieuwenhuizen, In: "Beyond the Quantum", eds. Th.M. Nieuwenhuizen et al., (World Scientific, Singapore) 2007, pp.332-342.
 - [27] A. Burinskii, Grav. Cosmol. **14**, 109 (2008) [arXiv:hep-th/0507109].
 - [28] H. B. Nielsen and P. Olesen, Nucl. Phys. B **61**, 45 (1973).
 - [29] M. S. Volkov and E. Wöhrner, Phys. Rev. D **66**, 085003 (2002) [arXiv:hep-th/0205157].
 - [30] N. Graham, Phys. Rev. Lett. **98**, 101801 (2007) [Erratum-ibid. **98**, 189904 (2007)] [arXiv:hep-th/0610267].
 - [31] F.E.Schunck and E.W.Mielke, in *Relativity and Scientific Computing*, edited by F.Hehl et.al., Springer, Berlin, 1966, pp.138-151.
S. Yoshida and Y. Eriguchi, Phys. Rev. D **56**, 762 (1997).
 - [32] V. Hamity, Phys. Lett. A **56**, 77 (1976). Phys. Rev. **143**, 1011 (19yy).
 - [33] P. A. M. Dirac, Proc. R. Soc. A **268**, 57 (1962).
 - [34] E. Witten, Nucl. Phys. B **249**, 557 (1985).
 - [35] J. R. Morris, Phys. Rev. D **53**, 2078 (1996) [arXiv:hep-ph/9511293].
 - [36] A. Burinskii, J. Phys. A **39**, 6209 (2006) [arXiv:gr-qc/0606097].
 - [37] W. Israel, Phys. Rev. D **15**, 935 (1977).
 - [38] A. Burinskii, in: Proc. of the XXV Int. Workshop on HEP and FT, (Protvino, IHEP 2002), pp.263-277, [arXiv:1006.1274].

- [39] A. Burinskii, Int. J. Mod. Phys. A **17**, 920 (2002) [arXiv:hep-th/0205127].
- [40] J. Ipser and P. Sikive, Phys. Rev. D **30**, 712 (1984).
- [41] M. Cvetič, S. Griffies, and H. H. Soleng, Phys. Rev. D **48**, 2613 (1993).
- [42] G. W. Gibbons, Commun. Math. Phys. **45**, 191 (1975).
- [43] A. H. Compton, Phys. Rev. **14**, 247 (1919).
- [44] The based on NED regularized KN solutions were presented in [18], while our attempts to get such solutions in the NED theory with two invariants ([17]) indicated their inconsistency with the GG form of metric (unpublished).
- [45] Recently, this twosheetedness found new holographic interpretation [8].
- [46] For $a = 0$, the Lópes source turns into the Dirac model of spherical electron [33].
- [47] In particular, for the parameters of an electron (in the units $\hbar = c = G = 1$) the bubble forms a highly oblate disk with the radius about the Compton length $a = J/m = 1/2m$ and the thickness equal to the "electromagnetic" radius of the electron, $r_0 = r_e = e^2/2m$. The degree of oblateness is $r_0/a = e^2 \sim 137^{-1}$.
- [48] It could be experiments related to very soft scattering of the polarized electrons. Note that Compton discussed experimental confirmation of their disklike shape in [43].